



McAuliffe, GA., Takahashi, T., Orr, R.J., Harris, P., & Lee, MRF.  
(2018). Distributions of emissions intensity for individual beef cattle  
reared on pasture-based production systems. *Journal of Cleaner  
Production*, 171, 1672-1680.  
<https://doi.org/10.1016/j.jclepro.2017.10.113>

Publisher's PDF, also known as Version of record

License (if available):  
CC BY

Link to published version (if available):  
[10.1016/j.jclepro.2017.10.113](https://doi.org/10.1016/j.jclepro.2017.10.113)

[Link to publication record in Explore Bristol Research](#)  
PDF-document

This is the final published version of the article (version of record). It first appeared online via Elsevier at <http://www.sciencedirect.com/science/article/pii/S0959652617324125> . Please refer to any applicable terms of use of the publisher.

## University of Bristol - Explore Bristol Research

### General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:  
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

## **Distributions of emissions intensity for individual beef cattle reared on pasture-based production systems**

**G.A. McAuliffe<sup>1,2</sup>, T. Takahashi<sup>1,2\*</sup>, R.J. Orr<sup>1</sup>, P. Harris<sup>1</sup> and M.R.F. Lee<sup>1,2</sup>**

<sup>1</sup> Rothamsted Research, North Wyke, Okehampton, Devon, EX20 2SB, UK,

<sup>2</sup> University of Bristol, Langford House, Langford, Somerset, BS40 5DU, UK

\* Corresponding author. Tel: +44 1837 883558. Email: taro.takahashi@rothamsted.ac.uk

### **Supplementary material (6 pages)**

Figure S1 Map of the North Wyke Farm Platform in Devon, UK

Figure S2 Distribution of emissions intensity per animal under each system

Table S1 Farm activities carried out under each system

Table S2 Distributions of uncertainty parameters assumed in Monte Carlo simulations

Table S3 Comparison of emissions intensity (kg CO<sub>2</sub>-eq/kg LWG) derived under two methods




Figure S1. Map of the North Wyke Farm Platform in Devon, UK. Green = permanent pasture (PP); Blue = white clover/high sugar grass (WC); Red = high sugar grass monoculture (HS).




Figure S2. Distribution of emissions intensity per animal under each system. Outliers located further than 1.5 times the interquartile range beyond the quartiles are each denoted with a cross (x).

*Table S1. Farm activities carried out under each system.*

Activity	PP	WC	HS
Ploughing	✗	✓	✓
Rolling	✓	✓	✓
Harrowing	✗	✓	✓
Seeding	✗	✓	✓
Fertiliser spreading	✓	✓	✓
Herbicide spraying	✗	✓	✓
FYM spreading (solid)	✓	✓	✓
Liming	✗	✓	✓
Mowing	✓	✓	✓
Silage making	✓	✓	✓

Table S2. Distributions of uncertainty parameters assumed in Monte Carlo simulations.

Emission source	Uncertainty	Distribution	Reference
<i>Animal/housing</i>			IPCC (2006)
Methane (EF and MM)	$\pm 20 \%$	Triangular	
Nitrous oxide (direct MM)	$SD^2 = 2$	Lognormal	
Nitrous oxide (indirect MM leaching)	-1500%/333%	Triangular	
Nitrous oxide (indirect MM volatilisation)	$SD^2 = 5$	Lognormal	
<i>Pasture</i>			IPCC (2006)
Nitrous oxide (direct)	$SD^2 = 3$	Lognormal	
Nitrous oxide (indirect leaching)	-1500%/333%	Triangular	
Nitrous oxide (indirect volatilisation)	$SD^2 = 5$	Lognormal	
Carbon dioxide (lime)	-50%/0%	Triangular	

*Table S3. Comparison of emissions intensity (kg CO<sub>2</sub>-eq/kg LWG) derived under two methods*

System	Representative animal approach				Individual animal approach			
	Mean	LL <sup>1</sup>	UL <sup>1</sup>	Range	Mean	Min	Max	Range
PP	17.8	15.0	21.5	6.5	18.4	16.3	21.7	5.4
WC	14.4	12.7	16.2	3.5	16.0	13.7	20.9	7.2
HS	19.0	16.3	22.5	6.2	20.2	16.6	25.6	9.0

<sup>1</sup> Lower and upper limit values of the 95% confidence interval estimated by Monte Carlo simulations.